

Stepwise formation of the bacterial flagellar system

Renyi Liu* and Howard Ochman*†‡

Departments of *Biochemistry and Molecular Biophysics and †Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721

Edited by Francisco J. Ayala, University of California, Irvine, CA, and approved March 8, 2007 (received for review January 11, 2007)

Elucidating the origins of complex biological structures has been one of the major challenges of evolutionary studies. The bacterial flagellum is a primary example of a complex apparatus whose origins and evolutionary history have proven difficult to reconstruct. The gene clusters encoding the components of the flagellum can include >50 genes, but these clusters vary greatly in their numbers and contents among bacterial phyla. To investigate how this diversity arose, we identified all homologs of all flagellar proteins encoded in the complete genome sequences of 41 flagellated species from 11 bacterial phyla. Based on the phylogenetic occurrence and histories of each of these proteins, we could distinguish an ancient core set of 24 structural genes that were present in the common ancestor to all Bacteria. Within a genome, many of these core genes show sequence similarity only to other flagellar core genes, indicating that they were derived from one another, and the relationships among these genes suggest the probable order in which the structural components of the bacterial flagellum arose. These results show that core components of the bacterial flagellum originated through the successive duplication and modification of a few, or perhaps even a single, precursor gene.

bacterial evolution | biological complexity | gene duplication

Bacterial flagella are complex and well honed organelles that provide swimming and swarming motilities and also play a central role in adhesion, biofilm formation, and host invasion (1). In the past several decades, extensive knowledge has accumulated about the structure, genetics, assembly, and regulation of flagella in widely diverse bacterial lineages (2–7). The typical bacterial flagellum consists of six components: a basal body (including MS ring, P ring, and L ring), a motor, a switch, a hook, a filament, and an export apparatus (2). In the best studied systems, those of *Escherichia coli* and *Salmonella enterica* sv. Typhimurium, >50 genes are involved in flagellar biosynthesis and function (3). Approximately half of these genes encode the structural components of the flagellum, and the rest are responsible for either the regulation of flagellar assembly or the detection and processing of environmental signals to which flagella respond.

Whereas *E. coli* and *Salmonella* have long served as the model organisms for studying flagellar assembly (2), there is extensive diversity among bacteria in the contents and organization of the gene complexes that specify flagella as well as structural variation in the flagellum itself (8, 9). For example, in Spirochaetes, flagella are located in the periplasm between the outer membrane sheath and cell cylinder (10); and, in accordance with their location, they have an enlarged C ring and rotor, and have a shape different from that seen in *Salmonella* (11). Furthermore, some bacteria, such as *Vibrio parahaemolyticus*, possess two flagellar systems (polar and lateral) that are encoded by distinct set of genes and use different motive forces (sodium and proton) but share a chemotaxis signal transduction system (12).

The bacterial flagellum has received attention as an exemplum of biological complexity; however, how this complexity and diversification have been achieved remains rather poorly understood. Although several scenarios have been posited to explain how this organelle might have been originated (13), the actual series of evolutionary events that have given rise to the flagellum,

as might be inferred from the relationships of all genes that contribute to the formation and expression of this organelle across taxa, has never been accomplished.

Insights into the evolution of the bacterial flagellum have been gained from the homologies between flagellar proteins and those functioning in other systems (13). For example, the sequence similarity between flagellum-specific ATPase FliI and the β -subunit of ATP synthase led to the speculation that flagellum possibly evolved from this highly conserved, membrane-bound enzyme, whose subunits rotate during catalysis of ATP from ADP (14). Because the flagellar motor proteins MotA/B are homologous to the motor proteins in the Tol-pal and TonB systems (15), the flagellum was hypothesized to have originated as a simple proton-driven secretion system (16). Most significantly, there are well established sequence and structural homologies between bacterial flagella and the type III secretion system (TTSS) demonstrating that the two apparatus derive from a common ancestor (17). Most evidence, including their much broader phylogenetic distribution, supports the view that the flagellum arose much earlier than the TTSS, which are largely limited to Proteobacteria (18–20).

Here, we take advantage of complete genome sequence data to trace the history of each gene involved in the assembly and regulation of the bacterial flagellum. Our results show that flagellum originated very early, before the diversification of contemporary bacterial phyla, and evolved in a stepwise fashion through a series of gene duplication, loss and transfer events. In this article, we focus on the evolution of the core set of flagellar genes that is uniformly present in all flagellated bacteria. The later evolving and lineage-specific components of the flagellar gene complexes remain to be addressed.

Results

Defining the Core Set of Flagellar Genes. By querying the genomes of flagellated bacteria for which complete genome sequences are available, we obtained the phylogenetic distribution of every gene known to be involved in the biosynthesis and regulation of flagella. To investigate the origin and evolution of the bacterial flagellar system, we then applied a phylogenetic profiling method (21) to assort genes into functional groups based on their co-occurrence and shared distributions across genomes. Genes with different functional roles have distinct phylogenetic distributions and profiles; however, most of genes whose protein products constitute the structural components of the flagellum are present in all bacterial phyla considered (Fig. 1). This distribution suggests this core set of structural genes originated before the divergence of the major bacterial lineages and includes 21 genes that specify proteins that form the filament (*fliC*,

Author contributions: R.L. and H.O. designed research; R.L. performed research; R.L. and H.O. analyzed data; and R.L. and H.O. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Abbreviation: TTSS, type III secretion system.

†To whom correspondence should be addressed. E-mail: hochman@email.arizona.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0700266104/DC1.

© 2007 by The National Academy of Sciences of the USA

hook length control gene *fliK*), have highly variable distributions and are excluded from the core set, even though some of the genes are known to be essential for proper functioning of the flagellar system in a particular species. (The evolutionary histories of these regulatory genes, along with that of a second bacterial flagellar system remain to be described.)

Phylogenetic Analysis of Flagellar Core Genes. To ascertain whether the 24 genes that form the flagellar core set have congruent evolutionary histories with one another, we compared the phylogenetic tree inferred for each core gene to that based on concatenated alignments of proteins encoded by 14 of the core genes. (These 14 genes were selected because they were present in all species included in this study and encoded the proteins having a high proportion of alignable positions.) For each of the 24 genes, all branches with >75% bootstrap values agreed with those in the concatenated tree, indicating that no alternative branching orders show strong support and that each of these genes has followed a common history in bacteria since they originated.

Congruence of Flagellar Genes with Organismal Phylogeny of Bacteria. The distribution of the 24 core genes among divergent bacterial phyla is most consistent with an ancient origin, pre-dating the shared ancestor of Bacteria. However, the distribution could have been achieved through later horizontal transfer. We tested these alternatives by comparing the phylogeny of the flagellar core proteins with the phylogeny of the corresponding bacterial phyla based on 25 universally distributed genes. The phylogenies are largely congruent on branches that have >75% bootstrap support; however, there are two inconsistencies between the core-gene and the organismal phylogenies; in the placement of both the alphaproteobacterial *Zymomonas mobilis* and a clade of three Betaproteobacteria within the Gammaproteobacteria (Fig. 2). Because individual flagellar genes within the core set show the same evolutionary history (see above), these incongruities have likely resulted from the transfer of the entire flagellar gene complexes between proteobacterial lineages after their separation from other major bacteria groups.

Core Flagellar Proteins Arose Through the Duplication and Diversification of a Single Precursor. When each of the 24 core flagellar proteins of *E. coli* are compared (via BLAST) to all proteins encoded in the *E. coli* genome, their best and often only hits are to other core flagellar proteins. Pair-wise comparisons among these core proteins revealed that ten are homologous to other core proteins when applying an *e*-value cutoff of 10^{-4} (Fig. 3). This pattern indicates that the structural genes specifying the portion of flagellum residing outside of cytoplasmic membrane (i.e., the rod, hook, and filament) are paralogs and were derived from one another through duplications.

Aside from these matches to other core proteins, pairwise comparisons of these flagellar proteins to the >4,000 nonflagellar proteins encoded by the entire *E. coli* genome recovered cumulatively a total of only 24 hits that reached the same level of significance. Among these matches, half (including some with *e*-values as low as $3e^{-10}$ to the flagellar core proteins) are involved in other secretion systems, such as the P pilus and the Type V secretion system, which is consistent with the idea that the flagellum originated as a secretion system. An additional 10 of the 24 hits (with *e*-values ranging from 10^{-5} to 10^{-6}) are membrane proteins, and the remaining two are prophage tail-fiber proteins. Thus, we conclude that despite their antiquity, the similarities among core proteins to one another are more common and, on average, stronger than to nonflagellar proteins.

Because the genes that constitute the core set are ancient and highly diverged, it is possible that some of the relationships among genes might not be recognized from analyses limited to

the *E. coli* flagellar complex. We repeated this analysis and compared the core gene set of each other flagellated bacterium to all proteins encoded in the corresponding genomes and among themselves, and we obtained a similar result, i.e., the best (and often the only) hits of the flagellar core genes were to other flagellar core genes. However, by extending this analysis beyond *E. coli*, the similarity-relationships and links among several other core genes were resolved. For example, a highly significant match between *fliM* and *fliN* (that was not detected for *E. coli* homologs) was evident in 15 genomes from diverse bacterial subdivisions (Fig. 3). In addition, the interacting export components encoded by *fliP*, *fliR*, and *fliQ* are related based on their protein sequences within several taxa. And even among the 10 *E. coli* core genes that originally showed similarity to one another, there were several new interconnections (e.g., *flgB* to both *flgE* and *flgG*, and between *flgL* and *flgK*) revealed by performing the analysis on other genomes. Cumulatively, each of the 24 core genes shows significant similarity to one or more of the other core genes (Fig. 3), a pattern that would result from their successive origination from one another by independent gene duplications and/or gene fusions.

The similarity among the proximal rod protein FlgF, the distal rod protein FlgG, and the hook protein FlgE exemplifies the relationships among these flagellar proteins (Fig. 4). FlgF and FlgG are of similar size (251 aa vs. 260 aa in *E. coli*) and show 31% amino acid identity over their entire lengths. In contrast, the *flgE* gene is much longer and appears to have evolved from *flgG* through an intragenic duplication that added a 160-aa domain to the N terminus of its encoded protein. PSI-BLAST searches reveal two significant alignments between FlgE and FlgG in *E. coli*: one with 24% identity between whole length of FlgG and the C terminus of FlgE (156–401 aa), and the other with 29% identity between the N terminus of two proteins (≈ 160 aa). That *flgE* evolved by a duplication is also supported by the fact that there are two versions of *flgE* in the genus *Bacillus*: among sequenced genomes, four species (*B. subtilis*, *B. clausii*, *B. licheniformis*, and *B. halodurans*) contain a shorter version, which is similar in length to *flgG*, and three species (*B. thuringiensis*, *B. cereus*, and *B. anthracis*) have the longer version.

From the matrix of relationships and protein sequence alignments of the flagellar core genes of *E. coli*, it is also possible to infer the order in which many of these genes and their corresponding structures originated. The low levels of protein identity among these paralogs, paralogous pairs are between 18% and 32% identical, required that we apply a method that combines the output of series of multiple alignment programs to derive a consensus alignment. The alignments on the terminal regions of the proteins, especially at the C terminus, offer the highest confidence. An unrooted neighbor-joining tree and a maximum-likelihood tree [supporting information (SI) Fig. 5] show that the rod proteins originated with either FlgB or FlgC, which are both short proteins, and then generated FlgF and FlgG (and hook protein FlgE) through a series of duplication events. The evolutionary relationships of these flagellar genes parallel the locations of their encoded proteins in contemporary flagella. The proximal, then distal, rod proteins precede (both evolutionarily and physically) the hook proteins, which preceded the hook-filament junction and filament proteins.

Discussion

Comparisons of the complete genome sequences of flagellated bacteria revealed that the flagellum is based on an ancestral set of 24 core genes for which homologs are present in genomes of all bacterial phyla. The most striking finding from our analysis is that these core genes originated from one another through a series of duplications, an inference based on the fact that they still retain significant sequence homology. The individual core genes show phylogenetic histories congruent with one another,

Phylogenetic Profiling. *E. coli* flagellar proteins are used as queries to search annotated proteins from complete genomes with BLASTP. Reciprocal best hits recovered at a cutoff of 10^{-5} among proteins in the *E. coli* and the queried genome were considered to be orthologs. If no ortholog was recovered, we first used TBLASTN to query the complete genome to confirm that its absence was not attributable to annotation errors. As a secondary check, PSI-BLAST was also used to find potential homologs. Homologs recognized from either the TBLASTN or PSI-BLAST searches were then examined for gene context. In cases where genes were in a colinear region or had at least one of the same neighboring genes as in *E. coli*, we regarded the sequences as orthologous to the *E. coli* gene. If an ortholog was still not found, we considered that particular flagellar gene as absent from a genome. To search for any flagellar genes or proteins that might be highly diverged from their *E. coli* homologs and go undetected in previous searches, we repeated the entire analysis using the flagellar gene/proteins of *Bacillus subtilis* as queries. In cases where there were multiple copies of homologs of a particular gene, paralogy was established if they had the original *E. coli* query gene as their best hit when searching against all *E. coli* proteins. TTSS proteins, although homologous to flagellar proteins, are easily identified because they generally have lower similarity values and are not contained within the flagellar gene neighborhood.

To evaluate whether the identification and distribution of particular flagellar proteins might be confounded by enhanced rates of protein evolution, we computed the overall percent protein identity for each flagellar protein in *E. coli* to its ortholog in *Salmonella enterica* sv. Typhimurium. (*Salmonella* and *E. coli* are used because orthology is easily ascertained, and alignments cover the entire lengths of proteins.) Most pairs of orthologs were >70% identical, and those with lower values (i.e., fast evolving) were FliC (54%), FliD (51%), FliK (47%), FliS (62%), and FliT (35%). Despite their relatively fast rates of evolution, FliC and FliD orthologs were detected in the genomes of all major bacterial groups and were included in the universally distributed core set. Moreover, this analysis also showed that “noncore” proteins are, by and large, not fast evolving and that their orthologs will be detected when present in a genome.

Similarity Among Core Proteins. To detect similarities among core proteins in each of 41 flagellated bacteria, each protein was compared with each of the other core proteins in the same genome by using the BL2seq program (with default options) in the NCBI BLAST package, applying an *e*-value cutoff of 10^{-4} .

Phylogenetic Analysis of Flagellar Proteins. Protein sequence alignments were generated in Muscle (34) with option (-maxiters 100), and poorly conserved regions were trimmed by using Gblocks method (35) with parameters (-b4 = 2 -b5 = n). PHYML (36) was used to construct maximum likelihood-based phylogenetic trees with 100 bootstrap replicates (with options 1 i 1 100 JTT e 4 e BIONJ y y). When multiple proteins from each genome were used to build a phylogenetic tree, individual protein alignments were first concatenated to form a single large alignment before phylogenetic analysis. To assess the evolutionary histories of individual members of the core set of flagellar genes, we compared the branching order inferred for alignments of each protein to that inferred for a concatenation of 14 well aligned proteins present in all flagellar systems. To construct a multiple sequence alignment of the highly divergent paralogous core proteins in *E. coli*, we used the *mcoffee* option of the T-Coffee program (37) (<http://igs-server.cnrs-mrs.fr/Tcoffee/tcoffee.cgi>).

Reconstruction of the Bacterial Species Tree. Based on the concatenated alignments of 31 single-copy genes (predominantly ribosomal proteins), Ciccarelli *et al.* (23) reconstructed the relationships for sequenced representatives of three domains of life. Of these 31 genes, 25 are present in all of the bacterial genomes that we considered, and we used the concatenated alignment of their encoded proteins to build the bacterial species tree.

We thank Nancy Moran for originally suggesting this project and for input on the manuscript; Eduardo Rocha, Vincent Daubin, and Emmanuelle Lerat for numerous helpful comments about the analyses; and Becky Nankivell for assistance in making figures. This work was supported by National Institutes of Health Grants GM56120 and GM74738 (to H.O.).

- Kirov SM (2003) *FEMS Microbiol Lett* 224:151–159.
- Macnab RM (2003) *Annu Rev Microbiol* 57:77–100.
- Macnab RM (2004) *Biochim Biophys Acta* 1694:207–217.
- Berg HC (2003) *Annu Rev Biochem* 72:19–54.
- McCarter LL (2006) *Curr Opin Microbiol* 9:180–186.
- Aldridge P, Hughes KT (2002) *Curr Opin Microbiol* 5:160–165.
- Soutourina OA, Bertin PN (2003) *FEMS Microbiol Rev* 27:505–523.
- Penn CW, Luke CJ (1992) *FEMS Microbiol Lett* 100:331–336.
- Bardy SL, Ng SYM, Jarrell KF (2003) *Microbiology* 149:295–304.
- Charon NW, Goldstein SF (2002) *Annu Rev Genet* 36:47–73.
- Murphy GE, Leadbetter JR, Jensen GJ (2006) *Nature* 442:1062–1064.
- McCarter LL (2004) *J Mol Microb Biotech* 7:18–29.
- Pallen MJ, Matzke NJ (2006) *Nat Rev Microbiol* 4:784–790.
- Rizzotti M (2000) *Early Evolution: From the Appearance of the First Cell to the First Modern Organisms* (Birkhauser, Boston).
- Cascales E, Llobes R, Sturgis JN (2001) *Mol Microbiol* 42:795–807.
- Musgrave I (2004) in *Why Intelligent Design Fails: A Scientific Critique of the New Creationism*, eds Young M, Edis T (Rutgers Univ Press, New Brunswick, NJ), pp 72–84.
- Blocker A, Komoriya K, Aizawa S (2003) *Proc Natl Acad Sci USA* 100:3027–3030.
- Hueck CJ (1998) *Microbiol Mol Biol Rev* 62:379–433.
- Nguyen L, Paulsen IT, Tchieu J, Hueck CJ, Saier MH (2000) *J Mol Microb Biotech* 2:125–144.
- Saier MH (2004) *Trends Microbiol* 12:113–115.
- Pellegrini M, Marcotte EM, Thompson MJ, Eisenberg D, Yeates TO (1999) *Proc Natl Acad Sci USA* 96:4285–4288.
- Daubin V, Gouy M, Perriere G (2002) *Genome Res* 12:1080–1090.
- Ciccarelli FD, Doerks T, von Mering C, Creevey CJ, Snel B, Bork P (2006) *Science* 311:1283–1287.
- Homma M, Derosier DJ, Macnab RM (1990) *J Mol Biol* 213:819–832.
- Pallen MJ, Penn CW, Chaudhuri RR (2005) *Trends Microbiol* 13:143–149.
- Minamino T, Gonzalez-Pedrajo B, Kihara M, Namba K, Macnab RM (2003) *J Bacteriol* 185:3983–3988.
- Horowitz NH (1945) *Proc Natl Acad Sci USA* 31:153–157.
- Jensen RA (1976) *Annu Rev Microbiol* 30:409–425.
- Fernald RD (2006) *Science* 313:1914–1918.
- Treize AEO, Collin SP (2005) *Curr Biol* 15:R794–R796.
- Olson EN (2006) *Science* 313:1922–1927.
- Jiang Y, Doolittle RF (2003) *Proc Natl Acad Sci USA* 100:7527–7532.
- Boone DR, Castenholz RW, Garrity GM (2001) *Bergey's Manual of Systematic Bacteriology* (Springer, New York).
- Edgar RC (2004) *Nucleic Acids Res* 32:1792–1797.
- Castresana J (2000) *Mol Biol Evol* 17:540–552.
- Guindon S, Gascuel O (2003) *Syst Biol* 52:696–704.
- Notredame C, Higgins DG, Heringa J (2000) *J Mol Biol* 302:205–217.

Corrections

BIOPHYSICS, CHEMISTRY. For the article “Destruction of long-range interactions by a single mutation in lysozyme,” by Ruhong Zhou, Maria Eleftheriou, Ajay K. Royyuru, and Bruce J. Berne, which appeared in issue 14, April 3, 2007, of *Proc Natl Acad Sci USA* (104:5824–5829; first published March 26, 2007; 10.1073/pnas.0701249104), the authors note that on page 5829, right column, first full paragraph, beginning on line 7, the sentence “Eight Cl⁻ counter ions were then added to neutralize the solvated system, giving a total system size of ≈40,000 atoms” should instead read: “Eight Cl⁻ counter ions for pH = 7 and 18 Cl⁻ counter ions for pH = 2 (ASP, GLU, and HIS residues are protonated), respectively, were then added to neutralize the solvated systems, giving a total system size of ≈40,000 atoms.” This error does not affect the conclusions of the article.

www.pnas.org/cgi/doi/10.1073/pnas.0705169104

EVOLUTION. For the article “Stepwise formation of the bacterial flagellar system,” by Renyi Liu and Howard Ochman, which appeared in issue 17, April 24, 2007, of *Proc Natl Acad Sci USA* (104:7116–7121; first published April 16, 2007; 10.1073/pnas.0700266104), the authors note that on page 7121, right column, line 4, the phrase “by using the BL2seq program (with default options)” should instead read: “by using the BL2seq program (with the ‘-F F’ option).” Additionally, in the Fig. 3 legend, line 3, the cutoff value 10⁻⁴ should instead appear as 9 × 10⁻⁴. Applying this cutoff yields a total of 70, not 24, hits of flagellar core proteins to nonflagellar proteins in the entire *E. coli* K-12 genome. These errors do not affect the conclusions of the article.

www.pnas.org/cgi/doi/10.1073/pnas.0705172104